

ver.4 このレシピのエネルギー		100	kcal	体重の0.75乗	1.81	kg	計算では	0.63	日分となります										
参考の表へ		1日に与えるエネルギー	158	kcal	体 重	2.2	kg	実際は	1.0	日分とする									
7/17(wed)		このレシピで摂取できる栄養				このレシピでの推奨量 *1				このレシピでの推奨量		このレシピでの推奨量		このレシピでの上限量 *2					
グラム						NRC				AAFCO				NRC				AAFCO	
水分		126.5				体重0.75乗ベース				乾燥重量ベース				体重0.75乗ベース				乾燥重量ベース	
乾燥重量		29.5																	
エネルギー		100.2																	
エネルギー密度		3.4 kcal/g																	
たんぱく質		6.6 22%				20%				10%		18%		15.0% ~ 23.0%				~	
内、動物性蛋白質		#VALUE! #####																	
脂 質		5.4 18%				11%				6%		5%		7.0% ~ 15.0%				~	
可溶性炭水化合物		13.0 44%																	
繊 不溶性		2.9 9.8%																	
維 総 量		4.3 14.5%																	
灰分		1.4 4.8%																	
ナトリウム		62 0.21%				0.16%				0.08%		0.06%		0.15% ~ 0.35%				0.07% ~ 0.25%	
カリウム		511 1.73%				0.86%				0.40%		0.60%							
カルシウム		125 0.42%				0.80%				0.40%		0.60%		0.50% ~ 1.00%				2.5%	
マグネシウム		39 0.131%				0.121%				0.060%		0.040%						0.3%	
リ ン		125 0.42%				0.61%				0.30%		0.50%		0.25% ~ 0.75%				0.20% ~ 0.30%	
カルシウム：リン比		1.00 : 1								1 : 1				1.2 : 1 ~ 1.4 : 1				~	

アミノ酸(参考)					
このレシピで摂取できる量			このレシピでの推奨量		
			NRC		AAFCO
	/DM	mg	体重0.75乗ベース	乾燥重量ベース	乾燥重量ベース
イソロイシン	1.0%	283	217	112	109
ロイシン	1.6%	471	397	201	174
リジン	1.5%	441	199	103	186
メチオニン	0.5%	150	199	97	
シスチン	0.4%	105			
メチオニン・シスチン計	0.9%	254	379	192	127
フェニルアラニン	0.9%	276	271	133	
チロシン	0.7%	213			
フェニルアラニン・チロシン計	1.7%	489	434	218	215
スレオニン	0.9%	264	253	127	142
トリプトファン	0.3%	82	83	41	47
バリン	1.1%	337	289	145	115
ヒスチジン	0.7%	205	112	56	53
アルギニン	1.2%	361	199	103	150

アミノ酸スコア(参考)						
このレシピで摂取できる量		窒素1gあたりのアミノ酸量	算定用評点ボタン		アミノ酸スコアを計算する(%)	
			1973年(一般用)	1985年(2-5歳)	1973年(一般用)	1985年(2-5歳)
(mg)						
イソロイシン	283	295	250	180	118	164
ロイシン	471	493	440	410	112	120
リジン	441	462	340	360	136	128
メチオニン・シスチン計	254	266	220	160	121	166
フェニルアラニン・チロシン計	489	511	380	390	135	131
スレオニン	264	276	250	210	110	131
トリプトファン	82	86	60	70	143	122
バリン	337	352	310	220	114	160
ヒスチジン	205	214	-	120	-	178
全窒素量(g)	1.0					

